

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 22-10-2007		2. REPORT TYPE Journal Article		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE  Ionic Liquids as Hypergolic Fuels (Postprint)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Tommy Hawkins & Ghanshyam Vaghjiani (AFRL/RZSP); Michael Rosander, Steven Chambreau, & Stefan Schneider (ERC); Gregory Drake (Redstone Arsenal)				5d. PROJECT NUMBER 23030423	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Air Force Research Laboratory (AFMC) AFRL/RZSP 10 E. Saturn Blvd. Edwards AFB CA 93524-7680				8. PERFORMING ORGANIZATION REPORT NUMBER  AFRL-RZ-ED-JA-2008-041	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  Air Force Research Laboratory (AFMC) AFRL/RZS 5 Pollux Drive Edwards AFB CA 93524-70448				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S NUMBER(S) AFRL-RZ-ED-JA-2008-041	
12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for public release; distribution unlimited (PA# 08079A)					
13. SUPPLEMENTARY NOTES Published in Energy & Fuels, 2008, 22, 2871–2872. (c) 2008 American Chemical Society.					
14. ABSTRACT After seminal work presented a decade ago, ionic liquids (IL) have now received a lot of attention as energetic materials for propellant applications. <sup>[1-3]</sup> In bipropellant rocket engines, it is desirable to achieve ignition by means of a hypergolic reaction and so to minimize system complexity. Hypergolic bipropellants are defined as fuel and oxidizer combinations that, upon contact, chemically react and release enough heat to spontaneously ignite, eliminating the need for an additional ignition source. This also makes them highly reliable for spacecraft and satellites which need to fire their rocket engines hundreds, or even thousands, of times during their lifetime. Unfortunately, no reliable <i>a priori</i> method for prediction of hypergolicity for fuel – oxidizer pairs is available today. The initial “hunting for the hypergol”, as John Clark entitled one of the chapters in his book <i>Ignition!</i> , took place mainly during WWII. <sup>[4]</sup> At that time, such toxic systems as “C-Stoff” (a mixture of N <sub>2</sub> H <sub>4</sub> ·H <sub>2</sub> O, methanol and water) and others consisting of triethyl amine, aniline, toluidine, xyldine and N-methyl aniline were developed. Today, environmental and health concerns are becoming more and more pressing in the propellant world. Nevertheless, hydrazine and its methylated derivatives are still the state-of-the-art fuels for bipropellant applications. Most of the problems handling hydrazine and its derivatives are related to their volatility, as they are carcinogenic vapor toxins. For these reasons, it is exceedingly attractive to replace hydrazine with ILs which have become paragons of environmental friendliness, green chemistry and low vapor toxicity.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Dr. Tommy Hawkins
Unclassified	Unclassified	Unclassified	SAR	3	19b. TELEPHONE NUMBER (include area code) N/A

## Ionic Liquids as Hypergolic Fuels

Stefan Schneider,<sup>\*,†</sup> Tommy Hawkins,<sup>†</sup> Michael Rosander,<sup>†</sup> Ghanshyam Vaghjiani,<sup>†</sup>  
Steven Chambreau,<sup>†</sup> and Gregory Drake<sup>‡</sup>

Air Force Research Laboratory, 10 East Saturn Boulevard, Building 8451, Edwards Air Force Base,  
California 93524, and Propulsion and Structures Directorate, United States Army Aviation and Missile  
Research, Development, and Engineering Center, AMSRD-AMR-PS-PT, Building 7120,  
Redstone Road Redstone Arsenal, Alabama 35898

Received April 24, 2008. Revised Manuscript Received June 2, 2008

After seminal work presented a decade ago, ionic liquids (IL) have now received a lot of attention as energetic materials for propellant applications.<sup>1–3</sup> In bipropellant rocket engines, it is desirable to achieve ignition by means of a hypergolic reaction and therefore to minimize system complexity. Hypergolic bipropellants are defined as fuel and oxidizer combinations that, upon contact, chemically react and release enough heat to spontaneously ignite, eliminating the need for an additional ignition source. This also makes them highly reliable for spacecraft and satellites, which need to fire their rocket engines hundreds or even thousands of times during their lifetime. Unfortunately, no reliable *a priori* method for prediction of hypergolicity for fuel–oxidizer pairs is available today. Here, we report the first ILs to manifest hypergolic ignition.

The initial “hunting for the hypergol”, as John Clark entitled one of the chapters in his book, *Ignition!*, took place mainly during World War II.<sup>4</sup> At that time, such toxic systems as “C-Stoff” (a mixture of N<sub>2</sub>H<sub>4</sub>·H<sub>2</sub>O, methanol, and water) and others consisting of triethyl amine, aniline, toluidine, xylydine, and *N*-methyl aniline were developed. Today, environmental and health concerns are becoming more and more pressing in the propellant world. Nevertheless, hydrazine and its methylated derivatives are still the state-of-the-art fuels for bipropellant applications. Most of the problems handling hydrazine and its derivatives are related to their volatility, because they are carcinogenic vapor toxins. For these reasons, it is exceedingly attractive to replace hydrazine with ILs, which are regarded as paragons of environmental friendliness, green chemistry, and low vapor toxicity.

Most of the focus in energetic ILs research has been on heterocyclic cations in combination with oxygenated anions, which frequently have stability and safety problems.<sup>5–7</sup> In a

bipropellant system, with ILs used as fuels, it is not necessary to use an oxygen-balanced IL. Furthermore, one can avoid oxygen-carrying anions completely, which, other things being equal, should make handling much safer because fuel and oxidizer are chemically separated. In a salt system, the positive charge of the cation makes it much more resistant toward oxidation. Consequently, electron- and fuel-rich anions are much easier to oxidize and thus would seem to hold the potential to promote hypergolic ignition.

Disappointingly, our first ignition tests carried out with ILs containing fuel-rich azide anions did not reveal any hypergolic properties.<sup>8</sup> However, the violent nature of their reactions was an encouragement to continue this line of work. In comparison, the same family of cations paired with the nitrate anion showed no noticeable reactivity toward inhibited, red-fuming nitric acid (IRFNA, ~83% HNO<sub>3</sub> plus 14% N<sub>2</sub>O<sub>4</sub> plus ~2% H<sub>2</sub>O plus 0.6% HF), white fuming nitric acid (WFNA, ~100% HNO<sub>3</sub>), or N<sub>2</sub>O<sub>4</sub>.

Therefore, the aim of this paper is to report on hypergolicity investigations of other fuel-rich anions, especially the dicyanamide anion. Dicyanamide was chosen not only because it is a fuel-rich anion but also because IL dicyanamides have some of the lowest viscosities among known ILs.<sup>9–11</sup> In bipropellant systems, pumps transport fuel and oxidizer, and therefore, excessive viscosity must be avoided.

Imidazolium-based ILs were first considered because they generally possess greater stability than their triazolium or tetrazolium analogues. Unsaturated sidechains, such as allyl, propargyl, and 2-butenyl were selected because it was hoped that they would initiate or promote hypergolic ignition. It has been demonstrated in the past that unsaturated, especially acetylenic, compounds show a tendency to hypergolicity and can even be used as fuel additives to promote hypergolic ignition.<sup>4</sup>

The starting materials, 1-*R*-3-methylimidazolium bromides (1–5), were prepared by alkylation of methylimidazole followed by metathesis with freshly prepared silver dicyanamide (Scheme 1). Because silver dicyanamide is insoluble in methanol, it can be used in excess, assuring complete conversion (see the Supporting Information).

(8) Schneider, S.; Hawkins, T.; Rosander, M.; Mills, J.; Vaghjiani, G.; Chambreau, S. *Inorg. Chem.* **2008**, in press.

(9) Yoshida, Y.; Muroi, K.; Otsuka, A.; Saito, G.; Takahashi, M.; Yoko, T. *Inorg. Chem.* **2004**, *43*, 1458.

(10) MacFarlane, D. R.; Golding, J.; Forsyth, S.; Forsyth, M.; Deacon, G. B. *Chem. Commun.* **2001**, 1430.

(11) Laus, G.; Bentivoglio, G.; Schottenberger, H.; Kahlenberg, V.; Kopacka, H.; Roeder, H.; Roeder, T.; Sixta, H. *Lenzinger Ber.* **2005**, *84*, 71.

(12) Gao, Y.; Gao, H.; Piekarski, C.; Shreeve, J. M. *Eur. J. Inorg. Chem.* **2007**, 4965.

\* To whom correspondence should be addressed. Fax: (+1) 661-275-5471. E-mail: stefan.schneider@edwards.af.mil.

<sup>†</sup> Air Force Research Laboratory.

<sup>‡</sup> U.S. Army Aviation and Missile RD&E Center.

(1) Hawkins, T. W.; Brand, A.; Petrie, M.; Bach, D.; McKay, M. Proceedings of the High Energy Density Matter, Contractors Conference, in Chantilly, VA, June 1–3, 1997; PL-TR-97-3057.

(2) Freemantle, M. *Chem. Eng. News* **2007**, *85*, 23.

(3) Shreeve, J. M.; Singh, R. P.; Verma, R. D.; Meshri, D. T. *Angew. Chem.* **2006**, *118*, 3664. Shreeve, J. M.; Singh, R. P.; Verma, R. D.; Meshri, D. T. *Angew. Chem., Int. Ed.* **2006**, *45*, 3584.

(4) Clark, J. D. *Ignition!*; Rutgers University Press: New Brunswick, NJ, 1972.

(5) Jones, C. B.; Haiges, R.; Schroer, T.; Christe, K. O. *Angew. Chem.* **2006**, *118*, 5303. Jones, C. B.; Haiges, R.; Schroer, T.; Christe, K. O. *Angew. Chem., Int. Ed.* **2006**, *45*, 4981.

(6) Hawkins, T.; Hall, L.; Tollison, K.; Brand, A.; McKay, M. *Propellants, Explos., Pyrotech.* **2006**, *31*, 196.

(7) Gálvez-Ruiz, J. C.; Holl, G.; Karaghiosoff, K.; Klapötke, T. M.; Löhnitz, K.; Mayer, P.; Nöth, H.; Polborn, K.; Rohbogner, J. C.; Suter, M.; Weigand, J. J. *Inorg. Chem.* **2005**, *44*, 4237.

Scheme 1




Table 1. Thermal and Physical Data of the Dicyanamide Salts 1–6

compound	$T_g$ (°C) <sup>a</sup>	$T_m$ (°C) <sup>b</sup>	$T_d$ (°C) <sup>c</sup>	$\eta$ (cP) <sup>d</sup> (25 °C)
1	−85		+207	42
2	−90		+210	27
3	−61	+17 <sup>e</sup>	+144	110
4		+49	+179	
5		+59	+184	
6	−66		+143	92

<sup>a</sup> Glass transition. <sup>b</sup> Melting point. <sup>c</sup> Decomposition onset. <sup>d</sup> Viscosity. <sup>e</sup> From ref 12.




Figure 1. ORTEP diagram showing connectivity, conformation, and the atom-numbering scheme for compound 5.

Physical data for known, 1 and 3, and novel, 2–5, compounds, are shown in Table 1 along with the 1-methyl-4-amino-1,2,4-triazolium analogue, 6, which is discussed later and was prepared by the same method.

Most substituted imidazolium dicyanamides are true room-temperature liquids. Compounds 4 and 5 are solid at ambient conditions, and compound 5 could be recrystallized from methanol layered with diethylether, giving crystals suitable for single-crystal X-ray determination. The structure analysis revealed an extensive hydrogen-bond network involving the dicyanamide anion (Figure 1). Similar to other ILs, solidification can be attributed to extensive hydrogen bonding.

As a simple test for hypergolicity similar to those used historically,<sup>4</sup> droplet tests were carried out using a glass cuvette, which was filled with ca. 1 mL of IRFNA or WFNA. A syringe generating ca. 10  $\mu\text{L}$  droplets was used to dispense the IL fuel (Figure 2a). A high-speed camera recording 500 frames/s was used to determine the ignition delay (ID) times (Table 2) by counting the frames between the droplet first hitting the surface of the oxidizer (Figure 2b) and the sign of the first visible ignition (Figure 2c). After ignition, the hypergolic compounds undergo self-sustained combustion (Figure 2d).




Figure 2. (a–d) Selected frames from the high-speed video of the hypergolic reaction of compound 3 and WFNA.

Table 2. ID Times for Hypergolic ILs with IRFNA and WFNA

compound dicyanamide	IRFNA ID (ms) <sup>a</sup>	WFNA ID (ms)
1	625	43
2	670	<sup>b</sup>
3	170	15
6	<sup>b</sup>	31
1-butyl-3-methyl-imidazolium <sup>c</sup>	<sup>b</sup>	47
1-butyl-1-methyl-pyrrolidinium <sup>c</sup>	<sup>b</sup>	44
<i>n</i> -butyl-3-methylpyridinium <sup>c</sup>	<sup>b</sup>	37

<sup>a</sup> Ignition delay time. <sup>b</sup> Not measured. <sup>c</sup> Purchased from Merck.

By this test, compounds 1–3 were shown to be hypergolic. The solids 4 and 5 were not tested because of their unsuitability as liquid propellants. One example of an IL based on a triazole, compound 6, and three commercially available IL dicyanamides, 1-butyl-3-methyl-imidazolium, 1-butyl-1-methylpyrrolidinium, and *n*-butyl-3-methyl-pyridinium, were also tested with WFNA and found to be hypergolic as well (Table 2).

On the basis of these results and somewhat unexpectedly, unsaturated substituents do not seem to play a determinative role in the ignition process. Rather, it appears that the key to hypergolicity in these systems is the dicyanamide anion.<sup>13</sup> For a suitable hydrazine replacement, ID times of less than 5 ms are preferred. Aerosol-type fuel and oxidizer sprays are being examined and should lead to shorter ID times as well as better reflecting real-world engine conditions.

With the discovery of these IL hypergols, a new path for transitioning these materials into bipropellant applications seems clear. It is expected that these systems can now be fine-tuned for energy content, performance, and desirable physical properties and that they could replace state-of-the-art, highly toxic hydrazine and its derivatives.

**Acknowledgment.** Financial support by the Air Force Office of Scientific Research (Mike Berman) is gratefully acknowledged.

**Supporting Information Available:** Crystallographic data for compound 5 in CIF format and further experimental details. This material is available free of charge via the Internet at <http://pubs.acs.org>.

EF800286B

(13) Chambreau, S. D.; Schneider, S.; Rosander, M.; Hawkins, T.; Gallegos, C. G.; Pastewait, M. F.; Vaghjiani, G. L. Submitted for publication.